

YEAST

# ENARTISFERM Q ET



Polyvalent yeast that can be used for direct inoculation

	<p><b>ORGANOLEPTIC CHARACTERISTICS</b></p> <p>EnartisFerm Q ET is a polyvalent yeast suitable for the production of white, red and rosé wines. When used in white and rosé vinification, it enhances varietal character of grapes, synthesizing fruity aromas which are stable over time. Moderate production of glycerol positively contributes to the sensations of volume and softness.</p> <p>In red vinification, it intensifies the extraction of polyphenolic substances and produces wines with great tannic structure, but also balanced at the end of fermentation and suitable for a long-term ageing.</p> <p>Due to its intrinsic characteristics and the special conditions adopted during its production, EnartisFerm Q ET (EasyTech) can be inoculated directly into must without the need for rehydration. Direct inoculation saves time and labor and facilitates yeast preparation, but above all, it reduces the risk of mistakes that can compromise a good fermentation process. When inoculated directly into juice, the high survival rate of EnartisFerm Q ET ensures a regular start to fermentation and its completion in the same time required by rehydrated yeast.</p>												
	<p><b>MICROBIOLOGICAL CHARACTERISTICS</b></p> <table border="0"> <tr> <td>Species</td> <td><i>Saccharomyces cerevisiae</i></td> </tr> <tr> <td>Fermentation temperature</td> <td>15 - 30°C (59-86°F)</td> </tr> <tr> <td>Lag phase</td> <td>medium</td> </tr> <tr> <td>Fermentation speed</td> <td>moderate</td> </tr> <tr> <td>Alcohol tolerance</td> <td>up to 16% v/v</td> </tr> <tr> <td>Killer factor</td> <td>neutral</td> </tr> </table>	Species	<i>Saccharomyces cerevisiae</i>	Fermentation temperature	15 - 30°C (59-86°F)	Lag phase	medium	Fermentation speed	moderate	Alcohol tolerance	up to 16% v/v	Killer factor	neutral
Species	<i>Saccharomyces cerevisiae</i>												
Fermentation temperature	15 - 30°C (59-86°F)												
Lag phase	medium												
Fermentation speed	moderate												
Alcohol tolerance	up to 16% v/v												
Killer factor	neutral												
	<p><b>ENOLOGICAL CHARACTERISTICS</b></p> <table border="0"> <tr> <td>Nitrogen needs</td> <td>medium</td> </tr> <tr> <td>Oxygen needs</td> <td>low</td> </tr> <tr> <td>Volatile acidity production</td> <td>low</td> </tr> <tr> <td>H<sub>2</sub>S production</td> <td>low</td> </tr> <tr> <td>SO<sub>2</sub> production</td> <td>medium-low</td> </tr> <tr> <td>Glycerol production</td> <td>medium</td> </tr> </table>	Nitrogen needs	medium	Oxygen needs	low	Volatile acidity production	low	H <sub>2</sub> S production	low	SO <sub>2</sub> production	medium-low	Glycerol production	medium
Nitrogen needs	medium												
Oxygen needs	low												
Volatile acidity production	low												
H <sub>2</sub> S production	low												
SO <sub>2</sub> production	medium-low												
Glycerol production	medium												
	<p><b>APPLICATIONS</b></p> <ul style="list-style-type: none"> <li>Production of young and medium aged white, red and rosé wines.</li> <li>Production of red wines for long-term ageing.</li> </ul>												
	<p><b>DOSAGE</b></p> <p>Primary fermentation: 20-40 g/hL (1.67 - 3.3 lb/1000 gal)</p> <p>The highest dosages are recommended in cases of compromised fruit, high sugar content and difficult microbiological conditions.</p>												
	<p><b>INSTRUCTIONS FOR USE</b></p> <p><i>Direct inoculation</i></p> <ul style="list-style-type: none"> <li>Pour yeast directly into juice or must (temperature &gt;15°C or 59°F) after or while filling the fermentation tank.</li> </ul> <p><b>Attention</b> - Alternatively, disperse the yeast in 10-20 parts of must (temperature &gt;15°C or 59°F) while stirring slowly to avoid clump formation, then add to the volume to be fermented.</p> <ul style="list-style-type: none"> <li>Pump-over to homogeneously distribute yeast in must.</li> </ul> <p><i>EnartisFerm Q ET can be used with rehydration, if desired.</i></p>												

The indications given here correspond to the current state of our knowledge and experience, however they do not relieve the user from compliance with safety and protection regulations or from improper use of the product.

	<p><i>With rehydration:</i></p> <ul style="list-style-type: none"> <li>▪ Dissolve dry yeast in 10 times its weight of clean, warm (35-40°C or 95-104°F) water. Stir gently.</li> <li>▪ Let suspension stand for 20 minutes, then gently stir again.</li> <li>▪ Add rehydrated yeast to juice or crushed grapes when you start to fill the fermentation tank. The difference in temperature between rehydrated yeast and juice should not exceed 10°C (18°F).</li> <li>▪ Homogenize by pump-over or mixing inoculated juice.</li> </ul>
	<p><b>PACKAGING AND STORAGE CONDITIONS</b></p> <p>10 kg</p> <p>Sealed package: store in a cool (preferably 5-15°C or 41-59°F) and dry area. Opened package: carefully reseal and store as indicated above; use quickly.</p>
	<p><b>COMPLIANCE</b></p> <p>The product is in compliance with: Codex Oenologique International.</p> <p>Product approved for winemaking in accordance with: Reg. (EU) 2019/934</p> <p><u>Product approved for winemaking by the TTB in accordance with 27 CFR 24.246.</u></p> <p>Legal Limit: N/A</p> <p>It contains E 491 Sorbitan monostearate</p>

The indications given here correspond to the current state of our knowledge and experience, however they do not relieve the user from compliance with safety and protection regulations or from improper use of the product.