

ENARTIS NEWS

THE IMPORTANCE OF BALANCED NUTRITION FOR YEAST HEALTH


IT'S NOT JUST ABOUT YAN

Appropriate, balanced nutrition is essential to guarantee a complete and regular alcoholic fermentation. Understanding and utilizing the balance between various amino acids and ammonia allows winemakers to improve the sensory profile of any wine, avoiding sluggish and stuck fermentations or sensory faults that will affect final wine quality.

Enartis has been researching all aspects that influence the correct development of yeast growth and, consequently, fermentation performance. This knowledge and research has given rise to an advanced set of tools.

YEAST CELL COMPOSITION

To understand the nutritional requirements of yeast for multiplication and proper metabolism, it is important to know yeast cell composition (*Figure 1*):



Of the total organic compounds in a yeast cell, 60-70% are nitrogenous substances, mainly structural proteins and enzymes. For this reason, it is important to measure the equivalent of metabolizable nitrogenous substances present in juice and adjust it accordingly. Nitrogen availability, regardless of the origin (amino acids or ammonia), will affect fermentation performance as well as the production of secondary metabolites and aromatic compounds during fermentation.

ESSENTIAL NUTRITIONAL REQUIREMENTS FOR YEAST

Balanced nutrition is essential for optimal status and biomass production, to improve yeast composition and performance, and produce desired aromatic compounds, avoiding the presence of off-flavors.

The contribution of nitrogenous substances is crucial for the physiological activity of yeast, among other microelements, vitamins, etc. Nitrogen can be assimilated from two different sources: amino acids or ammonia.

When **ammonia** (NH_4^+) is provided, yeast uses the nitrogen to synthesize proteins and enzymes. To produce amino acids from ammonia, yeast have to perform a long transformation process, **requiring large amounts of time and energy**. On the other hand, when **amino acids** are provided, yeast can store them for later use **without consuming energy** for their synthesis. Furthermore yeast can “decide” which metabolic pathways are more beneficial at that moment: to synthesize proteins, enzymes or other amino acids that will be used as nutritional sources to produce secondary products such as aromas (*Figure 2*).




Figure 2: Intracellular nitrogen pathways used depending on the nutritional source (amino acids or NH₄⁺).

Essential elements yeasts require for regular and complete fermentation are:

- Amino acids** – Naturally present in grapes in varying amounts depending on grape condition, variety, etc. Amino acids are the key compounds to provide balanced nutrition, guaranteeing optimal fermentation.

There are many types of amino acids, but not all of them are assimilated at the same speed by yeast (Table 1).

- Ammonia** – As shown in Table 1, it is not the element that yeast prefer since it is Class C. Of course, if the medium is deficient in yeast’s “favorite” amino acids, the yeast will consume ammonia first. The same occurs if a large amount of initial ammonia is provided.
- Vitamins** (biotin, thiamin, pantothenic acid, folic acid, etc.) and **microelements** (potassium, magnesium, phosphorus, sulfur, etc.) are considered essential growth factors for yeast. Therefore, it is important that these additions are made at the beginning of fermentation.
- Lipids** such as sterols and unsaturated fatty acids are considered survival factors as they are crucial for yeast membrane function. Yeast can also produce them by consuming oxygen. Lipid deficits in must leads to early cell death, fermentation problems, and increased volatile acidity.


Class A	Class B	Class C	Class D
Aspartate	Histidine	Alanine	Proline
Asparagine	Isoleucine	Ammonia	
Arginine	Leucine	Glycine	
Glutamate	Methionine	Phenylalanine	
Glutamine	Valine	Tryptophan	
Lysine		Tyrosine	
Serine			
Threonine			

Table 1: Classification of amino acids by time of consumption in must. Class A are the most preferred amino acids by yeast, and Class D cannot be assimilated by yeast.


MYTHS OF YEAST ASSIMILABLE NITROGEN (YAN)

In winemaking, there is a common misconception that a must needs to have 150-250 ppm of YAN to carry out a regular and complete alcoholic fermentation. It is often assumed that a higher dose of ammonia will help the development of yeast. Therefore, large quantities of diammonium phosphate (DAP) are added. Current literature and internal trials have shown that equivalent biomass production is attained when correct, balanced

nutrition is provided, regardless of the amount of YAN supplied (*Graph 1*).



Graph 1: Biomass production after 72h.




Graph 2: Comparison of different fermentation kinetics according to the nutrition source.

A must treated with 400 mg/L DAP and a must treated with 2 g/L autolyzed yeast contain a similar amount of yeast assimilable nitrogen. However, a significant difference in fermentation speed is observed (*Graph 2*). This is because the must treated with autolyzed yeast is rich in amino acids, microelements, vitamins and other compounds, not just ammonia.

All these elements are bio-active molecules preferred by yeast for multiplication and sugar metabolism - the main goal of fermentation. For this reason, it is important to consider the nutrient composition and the importance of the roles played by each component. This is what determines fermentation kinetics, not YAN.


ENARTIS NUTRIENT RANGE: CHARACTERISTICS AND APPLICATION TIME

By studying the nutritional needs of yeast, Enartis has developed the **Nutriform** range of nutrients to provide the most important elements at each stage of fermentation (*Graph 3*):




Graph 3: Application time of Nutriform range depending on the fermentation phase.

APPLICATION TIMING	NUTRIFERM RANGE	COMPOSITION CHARACTERISTICS	REASON TO USE
YEAST REHYDRATION or INOCULATION	NUTRIFERM AROM PLUS	<ul style="list-style-type: none"> • Rich in aromatic amino acid precursors (Graph 4) to promote the synthesis of higher alcohols that will later be converted into acetate esters: <ul style="list-style-type: none"> - Branched chain amino acids: valine, isoleucine, leucine. - Aromatic amino acids: tyrosine, phenylalanine, tryptophan. • Rich in essential growth factors such as vitamins and microelements to ensure optimal yeast function. 	<ul style="list-style-type: none"> • Promote the synthesis of secondary aromas. • Provide essential elements for yeast growth (amino acids, vitamins, microelements, etc.). • Improve yeast acclimatization to any condition. • Formulated to be added directly to juice without prior dissolving (Easytech).
	NUTRIFERM ULTRA	<ul style="list-style-type: none"> • Rich in Class A amino acids (Graph 5). • Abundant essential growth factors such as vitamins and microelements to ensure optimal yeast growth. • Undergoes special production process for immediate bio-availability. 	<ul style="list-style-type: none"> • Provides essential elements for yeast growth (amino acids, vitamins, microelements, etc.). • Stimulates protein synthesis, improving the physiological activity and longevity of yeast. • Improves yeast acclimatization to any condition. • Formulated to be added directly to juice without prior dissolving (Easytech).
1/3 ALCOHOLIC FERMENTATION	NUTRIFERM ADVANCE	Inactivated yeast with DAP and cellulose. Formulated to keep yeast functional until complete sugar depletion.	<ul style="list-style-type: none"> • Maintains the vital activity of yeast, helping the yeast to effectively finish fermentation. • Strengthens the cell walls of yeast. • Detoxifies the medium.
1/2 ALCOHOLIC FERMENTATION	NUTRIFERM NO STOP	Inactivated yeast, rich in survival factors (sterols, long-chain fatty acids, etc.), and yeast hulls that help detoxify juice.	<ul style="list-style-type: none"> • Regenerates the cell membrane and its fluidity to ensure successful alcoholic fermentation. • Detoxifies the medium by absorbing compounds that can inhibit fermentation, such as pesticide residues, medium-chain fatty acids, etc. • Recommended in difficult conditions to prevent or treat sluggish and/or stuck fermentations.



Graph 4: The addition of 20 g/hL Nutriferm Arom Plus increases the availability of aromatic amino acid precursors by 17% compared to the control (equivalent to any amount of DAP addition).



Graph 5: The addition of 20 g/hL Nutriferm Ultra increases the availability of the most important amino acids for yeast growth by 34% (Class A, Table 1) compared to the control (equivalent to any amount of DAP addition).

References: Jones. M.; Absorption of amino acids from wort by yeasts; in "Journal of the Institute of Brewing"; 1964.

Stay in touch with our newsletter.

SUBSCRIBE

www.enartis.com/en-au/newsletter